

Continuous and Discontinuous Phase Transitions in Quantitative Genetics: the role of stabilizing selective pressure

Annalisa Fierro^{1,2}, Sergio Coccozza³, Antonella Monticelli⁴, Giovanni Scala^{2,5}, and Gennaro Miele^{2,5}

¹ *CNR-SPIN Complesso Univ. Monte S. Angelo, Via Cinthia, I-80126 Naples, Italy*

² *Dipartimento di Fisica, Università degli Studi di Napoli "Federico II",
Complesso Univ. Monte S. Angelo, Via Cinthia, I-80126 Naples, Italy*

³ *Dipartimento di Medicina Molecolare e Biotecnologie Mediche,
Università degli Studi di Napoli "Federico II", Naples, Italy*

⁴ *Istituto di Endocrinologia ed Oncologia Sperimentale, CNR Napoli, Naples, Italy and*

⁵ *INFN - Sez. Napoli, Complesso Univ. Monte S. Angelo, Via Cinthia, I-80126 Naples, Italy*
(Dated: September 9, 2014)

By using the tools of statistical mechanics, we have analyzed the evolution of a population of N diploid hermaphrodites in random mating regime. The population evolves under the effect of drift, selective pressure in form of viability on an additive polygenic trait, and mutation. The analysis allows to determine a phase diagram in the plane of mutation rate and strength of selection. The involved pattern of phase transitions is characterized by a line of critical points for weak selective pressure (smaller than a threshold), whereas discontinuous phase transitions characterized by metastable hysteresis are observed for strong selective pressure. A finite size scaling analysis suggests the analogy between our system and the mean field Ising model for selective pressure approaching the threshold from weaker values. In this framework, the mutation rate, which allows the system to explore the accessible microscopic states, is the parameter controlling the transition from large heterozygosity (*disordered* phase) to small heterozygosity (*ordered* one).

I. INTRODUCTION

The need of applying a statistical approach arises in physics when the global properties of ensembles containing a huge number of elementary constituents are studied. In this case, the behavior of the single element (microstate) is irrelevant in favor of more informative averages on the whole ensemble (macrostate). In this sense, the situation is strictly analogous in both Population and Quantitative Genetics, where the analysis of allele frequencies and phenotype distribution parameters is preferred to the genetic/phenotypic description of each individual. The analogy between Quantitative Genetics and thermodynamics, noted from the very beginning by R. A. Fisher himself [1], allowed to introduce a concept of entropy for Population Genetics which satisfies the analogous of H-theorem [2]. Such information entropy measure ensures an exact solution at statistical equilibrium [2–5]. This point of view, can be alternatively seen by defining a free fitness, namely the entropy divided by population size plus the mean fitness. The free fitness is maximized at equilibrium when natural selection and drift (random sampling) are at work [2, 5], and provides an analogous of free energy in thermodynamics. For a summary of the correspondence between evolutionary dynamics and statistical physics see, for example, Table 1 of Ref. [4]. Following the above analogy, in Ref. [6] the authors study the evolution of a polygenic trait under stabilizing selection, mutation and genetic drift, following the evolution of the expectation of the trait mean, the genetic variance, and a measure of the expected heterozygosity. Interestingly, for population size N and mutation rate μ , such that $N\mu < 1/4$, the macroscopic quantities change their behavior showing a kind of *phase transition*. Moreover,

abrupt and gradual changes in the optimum, as well as changes in the strength of stabilizing selection seem to induce hysteresis phenomena, which can be understood imaging that the macroscopic trajectories keep memory of the underlying genetic states.

In the present paper, we analyze such phase transition phenomena looking for a more detailed statistical mechanics description. To this aim, we perform numerical simulations of a stochastic model, extensively analyzed in the past years (see for instance Ref. [3], and references therein). It consists in a population of N diploid hermaphrodite individuals, reproducing in pairs in a random mating regime, evolving under the effect of drift, selective pressure in form of viability, and mutation. Following Wright's seminal paper [7], we consider M different biallelic genes additively combining on the character, and the individual viability following a Gaussian profile in the trait. The model differs from those of Ref. [4] for the large interval of mutation rate considered here, and for the chosen fitness of the phenotype. Furthermore, in our analysis, the size of the population N plays the same role of the finite dimension in a physical system.

II. THE MODEL

We consider a system of N diploid individuals, sexually reproducing with random mating between any pairs of individuals. Each individual is represented by two sequences of M variables, σ_{kj} (where $k = 1, 2$ stands for the two genome replicas, and $j = 1, \dots, M$ runs on different loci). We refer to σ_{kj} as alleles, and assume that each allele can take two values, ± 1 . At each generation, all individuals are removed, and a new generation of same

size N is formed by off-springs of the previous individuals. A mutation rate, μ , is introduced, as the probability of an allele to mutate at each generation ($\sigma_{kj} \rightarrow -\sigma_{kj}$). Similar models were studied in Ref.s [8, 9].

We denote the additive polygenic phenotype random variable with $p = \sum_{j=1}^M \sum_{k=1}^2 \sigma_{kj}$. A stabilizing selective pressure on such a phenotype can be implemented via the survival probability of an offspring, typically known as *viability*,

$$S(p) \equiv \mathcal{N} \omega \exp \left[-(p - p_m)^2 \omega^2 / 2 \right], \quad (1)$$

where $\mathcal{N}$ is the suitable normalization constant, ω measures the *strength of selection*, and p_m stands for the optimum phenotype (hereafter, we choose for simplicity $p_m = 0$). For $\omega = 0$, there is no selective pressure, and all the microscopic states are equivalent. Whereas, for $\omega \rightarrow \infty$, the selective pressure is at maximum, and the only surviving individuals are those with $p = p_m$. Thus, in general, the effect of the selective pressure is to reduce the accessible phase space to those microscopic states satisfying *in probability* the constraint on the phenotype.

Let us denote with $\rho_{\text{in}}(j)$ the initial frequency of the allele “1” in locus j . We consider for simplicity a starting population where the M variants are ± 1 with equal probability, namely $\rho_{\text{in}}(j) = 0.5$, $\forall j = 1, \dots, M$. This ansatz does not restrict the generality of the analysis, in fact for finite populations, random drift rapidly homogenizes the initial distribution of allele frequencies, making the final results almost independent of the initial assumption on $\rho_{\text{in}}(j)$ [6].

To describe the macroscopic state of the system, we introduce the following quantity:

$$\bar{q} \equiv \frac{1}{M} \sum_{j=1}^M \langle |q(j)| \rangle, \quad (2)$$

with

$$q(j) \equiv \frac{1}{2N} \sum_{i=1}^N \sum_{k=1}^2 \sigma_{ik}(j), \quad (3)$$

where $\langle \dots \rangle$ stands for the average over ~ 30 initially identical realizations of the system (hereafter simply denoted by *ensemble* of populations). Note that the definition of $\bar{q}$ is inspired by a statistical mechanics analogue. In fact, if the quantity $q(j)$ of Eq. (3) resembles the magnetization per spin of an Ising model of $2N$ spins, $\bar{q}$ of Eq. (2) gives the average of the magnetization modulus once properly normalized [10]. Following the same analogy, we also introduce the *susceptibility*, as

$$\bar{\chi} \equiv \frac{1}{M} \sum_{j=1}^M \chi(j), \quad (4)$$

with $\chi(j) \equiv 2N (\langle q(j)^2 \rangle - \langle |q(j)| \rangle^2)$. It is interesting to note that the *magnetization* $q(j)$ of j -th locus is related to

the *expected* heterozygosity (fraction of heterozygous individuals expected on the basis of Hardy-Weinberg equilibrium condition) in the same locus, denoted by $h_s(j)$, which is a more familiar quantity in the Population Genetics context. Indeed, one can easily prove that

$$\begin{aligned} h_s(j) &\equiv \frac{1}{4N^2} \sum_{i,l=1}^N \sum_{k,n=1}^2 (1 - \delta_{\sigma_{ik}(j)\sigma_{ln}(j)}) = \\ &= \frac{1}{2} (1 - q(j)^2). \end{aligned} \quad (5)$$

It is worth observing, that in our case $h_s(j)$ essentially coincides with the *observed* heterozygosity (observed fraction of heterozygous individuals). From Eq. (5), we see that the minimum of magnetization corresponds to the maximum of heterozygosity, and vice-versa. Denoting with $\bar{H}_s$ the average of $h_s(j)$ over different loci and on the ensemble, one can easily prove that

$$\bar{H}_s = \frac{1}{2} \left(1 - \frac{\bar{\chi}}{2N} - \bar{q}^2 \right) \quad (6)$$

for a very large number of realizations. This occurs since, in this limit, $\langle |q(j)| \rangle$ is independent of j .

III. RESULTS

For any fixed set of parameters N , M , ω , and μ , we follow numerically the evolution of a given population till it asymptotically reaches a stationary state, which we refer to as *steady* state, where we evaluate $\bar{q}$ and $\bar{\chi}$. Let us start by focusing our attention on the role played by mutation rate and selection strength only, and to this aim we fix the values of N and M . In general, the system reaches the steady state for values of generation number, which depend on ω and μ . Two different behaviors are observed in the regime of small and large selective pressure strength, respectively.

In Fig. 1, we report $\bar{q}$ as a function of μ for different values of ω . As shown in figure, by decreasing the mutation rate μ , $\bar{q}$ goes from small- $\bar{q}$ (which vanishes in the limit of large- N) to $\bar{q} \sim 1$. Note that, for large μ , the alleles ± 1 have roughly equal probability (namely, $h_s(j) \sim 0.5$ for each locus), and hence the steady state does not significantly differ from the initial one. On the contrary, for small μ , in the steady state $h_s(j) \sim 0$ for each locus. In this case, the individuals are just clones, namely, for each realization, the population is represented by a unique genome that is a generic combination of ± 1 in a neighborhood of the phenotype optimum (exactly in the optimum for $\omega \rightarrow \infty$). The crossover from the state with small $\bar{q}$ (large heterozygosity) to the state with $\bar{q} \sim 1$ (small heterozygosity) is characterized by a maximum in the susceptibility, $\bar{\chi}$. The value of μ corresponding to such a maximum is a monotonic increasing function of ω . Moreover, concerning its dependence on M , it is interesting to observe that it simply scales




FIG. 1: Order parameter, $\bar{q}$ vs μ , in the steady states for $N = 1000$ and $\omega = 0, 0.1, 0.2, 0.4, 0.5, 1, \infty$ (from left to right). The continuous lines are guides for eyes.

as $1/M$, as one can expect since $2\mu M$, representing the mutation rate per individual, is the relevant quantity, ruling the mutations during evolution. From Fig. 1, it can be easily observed that for weak selection strength, roughly $\omega < 0.4$ (with blue circles in figure corresponding to $\omega = 0.4$), one has a smooth crossover that becomes abrupt for larger ω .

To better analyze the nature of these steady states, and the crossover from small- $\bar{q}$ states to large- $\bar{q}$ ones, we perform the following numerical experiment. For any value of ω , starting from a configuration at high mutation rate, we decrease μ at a given rate $\dot{\mu} \equiv \Delta\mu/\Delta n$ (n denoting the generation number). In other words, the system is kept at a given value of the mutation rate for an interval Δn , and, at the end of it, $\bar{q}$ and $\bar{\chi}$ are measured. Afterward, the value of μ is decreased of $\Delta\mu$ and the procedure is iterated till μ reaches zero. At this point the procedure is inverted and μ is increased at the same rate, in analogy to a physical system, first cooled and then heated at given rate. For any μ , two states are considered macroscopically equivalent if the measured values of $\bar{q}$ and $\bar{\chi}$ coincide, in analogy with a thermodynamical approach. This procedure confirms the presence of two different regimes, for $\omega < \omega_c$ and $\omega \geq \omega_c$, respectively, where the threshold $\omega_c \simeq 0.4 + \mathcal{O}(N^{-1})$.

In the region of small selective pressure, namely for $\omega < \omega_c$, plotting $\bar{q}$ as function of μ , for small enough values of the *cooling* rate, $\dot{\mu}$, we find that the curve does not depend on $\dot{\mu}$ at all. Moreover, the two branches, obtained by decreasing and increasing μ respectively, always coincide. In the main frame of Fig. 2, $\bar{q}$ vs μ is plotted for two different values of the cooling rate, and assuming $\omega = 0.1$. Similar behaviors are observed plotting $\bar{\chi}$ vs μ (see the inset of Fig. 2). Moreover, as it can be seen from Fig. 2, the states obtained with this




FIG. 2: **Main frame:** Order parameter, $\bar{q}$ vs μ , for $\omega = 0.1$ and $N = 1000$. The full lines are obtained first cooling the system at fixed $\dot{\mu}$ (the red line corresponds to 10^{-8} and the blue line to 10^{-9}), and then heating it at the same rate. The pink stars correspond to the steady states. **Inset:** Susceptibility, $\bar{\chi}$ vs μ , with the same symbols as in the main frame.

procedure coincide with the above defined steady states (pink stars in Fig. 2), and in some sense they can be considered *equilibrium* states of the system. In this region, the crossover from large to small heterozygosity, ruled by the mutation rate and characterized by a susceptibility maximum, strongly resembles a continuous (second order) transition in a physical system. In the following, we will study this transition as a usual critical phenomenon.

A. Fourth order cumulant and finite size scaling analysis at small selective pressure

From the intersection of the fourth order cumulant, evaluated for different sizes of the system, the critical mutation rate, $\mu_c \equiv \lim_{N \rightarrow \infty} \mu_c(N)$, is estimated, and a finite size scaling analysis is performed in order to evaluate the critical exponents of the transition.

For an Ising model with vanishing magnetic field, the reduced fourth order cumulant of the order parameter [11] is given by

$$U_4 = 1 - \frac{\langle m^4 \rangle}{3\langle m^2 \rangle^2}, \quad (7)$$

where m is the magnetization. Following Ref. [11], as the system size $N \rightarrow \infty$, $U_4 \rightarrow 0$ for $T > T_c$ and $U_4 \rightarrow 2/3$ for $T < T_c$. For large enough values of the size N , all curves representing U_4 as a function of temperature cross in a point whose location gives the critical point.

A natural extension of Eq. (7) to our system is

$$\overline{U}_4 \equiv \frac{1}{M} \sum_{j=1}^M U_4(j), \quad (8)$$

where

$$U_4(j) = 1 - \frac{\langle q(j)^4 \rangle}{3\langle q(j)^2 \rangle^2} \quad (9)$$

is the fourth order cumulant of j -th locus, and $\overline{U}_4$ denotes the average over the M loci.




FIG. 3: $\overline{U}_4$ vs μ , for $\omega = 0.2$ and different values of N . The crossing point is $\mu_c \sim 0.0021$. The continuous lines are guides for eyes.




FIG. 4: Critical mutation rate, μ_c (red circles) vs ω , compared with $\mu_c(N)$, for $N = 1000$ (blue stars).

In the present case, increasing the mutation rate, in each locus j it is observed a transition from a disordered




FIG. 5: $N^a \overline{q}$ (upper curves) and $N^{-b} \overline{\chi}$ (lower curves) vs ϵN^c , for $\omega = 0.2$ and different values of N . In this case, we have $a = 0.25$, $b = 0.6$, $c = 0.5$, $\epsilon = (\mu - \mu_c)/\mu_c$ and $\mu_c = 0.0021$.




FIG. 6: $N^a \overline{q}$ (upper curves) and $N^{-b} \overline{\chi}$ (lower curves) vs ϵN^c , or $\omega = 0.05$ and different values of N . In this case, we have $a = 0.15$, $b = 0.7$, $c = 0.5$, $\epsilon = (\mu - \mu_c)/\mu_c$ and $\mu_c = 0.00065$.

phase to an ordered one. This is in perfect analogy with the Ising model. Hence, we expect that varying the size N of the system, all curves for $\overline{U}_4$ as a function of μ cross in a point which provides the critical mutation rate $\mu_c \equiv \lim_{N \rightarrow \infty} \mu_c(N)$. Note that μ_c keeps a dependence on ω .

In Fig. 3, $\overline{U}_4$ is plotted as a function of μ for different values of the size N , having fixed $\omega = 0.2$. As expected, in the limit of large N we find that $\overline{U}_4$ tends to $2/3$ and to zero for small and large mutation rate, respectively. The crossing point, μ_c , represented by the red circles in Fig. 4, increases by increasing ω and vanishes in the limit $\omega \rightarrow 0$. Blue stars in the same figure correspond to the




FIG. 7: **Main frame:** $\bar{q}$ and $N^{-1}\bar{\chi}$ vs μN , for $\omega = 0$ and $N = 1000, 2000, 5000, 10000$. **Inset:** Critical value of the mutation rate, μ_c vs N . The full line is the power law function A/N , with $A \sim 0.1$.

maximum points of the susceptibility, namely $\mu_c(N)$ for $N = 1000$. We find that $\mu_c(N)$ is smaller than μ_c for each non vanishing ω , whereas this behavior reverses for $\omega = 0$.

Next, the critical behavior of the order parameter and of the susceptibility are studied. Extending the predictions from finite size scaling analysis in the Ising model [11] to the present system, we expect that near the critical point

$$\begin{aligned}\bar{q} &= N^{-a} q_0(\epsilon N^c), \\ \bar{\chi} &= N^b \chi_0(\epsilon N^c),\end{aligned}\quad (10)$$

where $\epsilon = (\mu - \mu_c)/\mu_c$, and q_0 and χ_0 are scaling functions. In Figs 5 and 6, $N^a \bar{q}$ and $N^{-b} \bar{\chi}$ are plotted for $\omega = 0.2$ and 0.05 respectively, as a function of ϵN^c , where the exponents a , b , and c are chosen in order to rescale the curves for different N onto a unique one. As it is clear from Figs 5 and 6 a departure from scaling laws Eq. (10), observed from $\mu < \mu_c$, is more evident for larger values of ω . The reason is that, increasing ω , for small mutation rate the system quickly becomes fully ordered (i.e., $\bar{q} \sim 1$), and the scaling does not hold.

In a d -dimensional physical system, a , b and c are related to the critical exponents, ν , β and γ , by the following relations [11]: $a = \beta/\nu d$, $b = \gamma/\nu d$, and $c = 1/\nu d$. Although the space dimension is here not defined at all, we can evaluate β and γ , from a , b and c , as $\beta = a/c$ and $\gamma = b/c$. In Table I, μ_c and the scaling exponents obtained for different ω are listed. Although the errors are rather large and further analysis is necessary, the critical exponents seem to change along the critical line, and to tend to the mean field Ising critical exponents (i.e., $\beta = 0.5$ and $\gamma = 1$) by approaching ω_c . We can speculate that the role of the euclidean dimension is here played

by the selective pressure, which controls the *energy* landscape of the system (the increasing of ω corresponds to an increasing of the energy barriers between two minima).

For $\omega = 0$, no crossing point is observed in the fourth order cumulant. Similar findings are obtained for $\omega = 0.01$. Consistently, the maximum point of the susceptibility $\mu_c(N)$ goes to zero as $1/N$ (see inset of Fig. 7) in the limit $N \rightarrow \infty$, and, as it is shown in main frame of Fig. 7, the following trivial finite size scaling is found:

$$\begin{aligned}\bar{q} &= q_0(\mu N), \\ \bar{\chi} &= N \chi_0(\mu N).\end{aligned}\quad (11)$$

It is worth observing, that due to this scaling behavior of $\bar{q}$ and $\bar{\chi}$, the heterozygosity, $\bar{H}_s$, defined in Eq. (6) results scale free for $\omega = 0$. Moreover, in absence of selective pressure, since the loci are independent, there is no dependence on M at all.

ω	$\mu_c(\pm 0.00005)$	a	b	c	β	γ
0	0	0	1	1	0	1
0.05	0.00065	0.15	0.7	0.5	0.3	1.4
0.1	0.0014	0.175	0.65	0.5	0.35	1.3
0.13	0.0017	0.2	0.6	0.5	0.4	1.2
0.2	0.0021	0.25	0.6	0.5	0.5	1.2

TABLE I: Critical mutation rate and scaling exponents, for different ω , are reported. Note that the scaling relation $2\beta + \gamma = \nu d$, which in terms of a and b becomes $2a + b = 1$, is almost everywhere verified. Note that the uncertainty on the a , b and c determination is ~ 0.05 , whereas the error on β and γ estimations are ~ 0.1 and ~ 0.2 , respectively.




FIG. 8: Order parameter, $\bar{q}$ vs μ , for $N = 1000$ and $\omega \rightarrow \infty$. The different cycles have been obtained by using different cooling rates. The black stars reproduce data plotted in Fig. 1 for $\omega \rightarrow \infty$, obtained by following the system up to $2 \cdot 10^6$ generation numbers.

B. Hysteresis cycles at large selective pressure

Interestingly, in the region of large selective pressure, for $\omega \geq \omega_c$, a metastable hysteresis appears between small values of $\bar{q}$ and 1, as shown in Fig. 8. Again, for large mutation rate, the system is at equilibrium in states with small- $\bar{q}$, and, for small mutation rate, is at equilibrium in states with $\bar{q} \sim 1$. However, the two branches, at small- $\bar{q}$ and $\bar{q} \sim 1$ respectively, are both observed for intermediate values of the mutation rate, depending on the pattern of μ -variation. Although by decreasing the cooling rate the hysteresis cycle shrinks, we always see two well distinct branches on our observation time scales. This behavior is reminiscent of a discontinuous (first order) transition, where metastable hysteresis is usually observed. In this case, the distinction of long-lived metastable states from equilibrium states is rather difficult, since the lifetime of the metastable states may be longer than the observation time. As it can be seen in Fig. 8, the states, obtained decreasing μ at small $\dot{\mu}$, coincide with the steady states, reached by the system for very large generation numbers. Hence, in this case the so-called steady states, which are stationary on our observation time scales, are likely metastable. Note that in Fig. 8 the hysteresis curves are plotted for $\omega \rightarrow \infty$, where this phenomenon is more evident.

Our findings are efficaciously summarized in Fig. 9, where the phase diagram for a system of $N = 1000$ individuals is shown in the plane $(\mu, 1/\omega)$. For $\omega < \omega_c$, we plot $1/\omega$ as function of the maximum point of $\bar{\chi}$, $\mu_c(\omega, N)$. The blue line should give, in the *thermodynamic* limit, a line of critical points, where the continuous transition from small- $\bar{q}$ to $\bar{q} = 1$ phase should be observed. Above ω_c , data depend on the cooling rate and the susceptibility displays two maxima, depending on the pattern of μ -variation. Red circles in Fig. 9 correspond to the maximum points of $\bar{\chi}$, along the hysteresis loop obtained at the smallest cooling rate, $\dot{\mu} = 10^{-9}$. In this region, the system behaves as a physical system undergoing a discontinuous transition controlled by the mutation rate.

C. Heterozygosities

The analysis on the dependence of the order parameter, $\bar{q}$, on the mutation rate, μ , is also carried out for the expected heterozygosity of the single population, $\bar{H}_s$, and for the expected heterozygosity measured on the set of populations as a whole, $\bar{H}_t$. As for $\bar{q}$, hysteresis cycles are observed for $\omega \geq \omega_c$ (see Fig. 11). Interestingly, $\bar{H}_s$ and $\bar{H}_t$ display different behaviors for small mutation rate. Figs 10 and 11 show that $\bar{H}_s \sim 0.5$ and $\bar{H}_t \sim 0.5$, for large mutation rate, whereas $\bar{H}_s \sim 0$ and $\bar{H}_t \sim 0.5$ (we expect $\bar{H}_t = 0.5$ for $N_p \rightarrow \infty$), for small mutation rate, where the independently evolving populations (although initially identical) reach fixation in generally different (but macroscopically equivalent) microstates, de-




FIG. 9: Phase diagram in the plane $(\mu, 1/\omega)$ for a system of size $N = 1000$ (see text for explanations). The continuous lines are guides for eyes.

veloping a genetic diversity. This phenomenon is in some sense analogous of the spontaneously symmetry breaking in physical system.




FIG. 10: **Main frame:** $\bar{H}_t$ vs μ , in the steady states for $N = 1000$ and $\omega = 0, 0.1, 0.2$ (from left to right). **Inset:** $\bar{H}_s$ vs μ , in the steady states for $N = 1000$ and $\omega = 0, 0.1, 0.2$ (from left to right). The continuous lines are guides for eyes.

IV. CONCLUSIONS

In summary, in this paper we have analyzed the evolution of a population of N diploid individuals, sexually reproducing with random mating, evolving under the effect of a Gaussian viability depending on an additive polygenic trait. Using the standard tools of Statistical




FIG. 11: Heterozygosities, $\overline{H}_s$ (red circles) and $\overline{H}_t$ (black circles) vs μ , for $N = 1000$ and $\omega \rightarrow \infty$, obtained first cooling the system and then heating it at given rate. Here $\dot{\mu} = 10^{-9}$

Mechanics, we show that the system displays a complex phase diagram with a transition from a disordered to an ordered phase, controlled by the mutation rate. We provide the phase diagram in the (μ, ω) plane, showing that the order of the transition changes depending on the strength of selection, being continuous for weak selective pressures and discontinuous for strong ones.

Acknowledgments The authors would like to thank L. Peliti and A. Coniglio for valuable discussions.

-
- [1] R.A. Fisher, in *The Genetical Theory of Natural Selection*, (Clarendon, Oxford, 1930).
 - [2] Y. Iwasa, J. Theor. Biol. **135**, 265 (1988).
 - [3] N.H. Barton, Genetical Research **54** 59 (1989).
 - [4] G. Sella and A.E. Hirsh, Proc. Natl. Acad. Sci. U.S.A., **102**, 9541 (2005).
 - [5] N.H. Barton, J.B. Coe, Jour. of Theor. Biology, **259**, 317 (2009).
 - [6] H.P. de Vladar, N.H. Barton, J R Soc Interface, **8**, 720 (2011).
 - [7] S. Wright, Genetics **16**, 16 (1931).
 - [8] M. Serva and L. Peliti, J. Phys. A: Math. Gen. **24** (1991) L705.
 - [9] P.G. Higgs and B. Derrida, J. Phys. A: Math. Gen. **24** (1991) L985.
 - [10] For vanishing magnetic field ($H = 0$) the Ising model undergoes a second order phase transition from a disordered paramagnetic phase (vanishing magnetization) to an ordered ferromagnetic one (not vanishing magnetization), at temperature T_c . In the ferromagnetic phase ($T \leq T_c$), the magnetization $\rightarrow 0$ at the critical temperature as a power law with exponent β . For fixed temperature $T < T_c$, a first order transition controlled by the magnetic field, is found for vanishing H .
 - [11] K. Binder Z. Phys. B Condensed Matter 1981, **43** 119; D.L. Landau and K. Binder, in *A Guide to Monte-Carlo Simulations in Statistical Physics*, (Cambridge University Press 2013).